

A POLYNOMIAL WEYL INVARIANT SPINNING MEMBRANE ACTION

Carlos Castro

Center for Theoretical Studies of Physical Systems
Clark Atlanta University, Atlanta, GA. 30314
November 2002

Abstract

A review of the construction of a Weyl-invariant spinning-membrane action that is *polynomial* in the fields, without a cosmological constant term, comprised of quadratic and quartic-derivative terms, and where supersymmetry is linearly realized, is presented. The action is invariant under a *modified* supersymmetry transformation law which is derived from a new $Q + K + S$ sum-rule based on the $3D$ -superconformal algebra .

A satisfactory spinning membrane Lagrangian has not been constructed yet, as far as we know. Satisfactory in the sense that a suitable action must be one which is *polynomial* in the fields, without (curvature) R terms which interfere with the algebraic elimination of the three-metric, and also where supersymmetry is linearly realized in the space of physical fields.

Lindstrom and Rocek [1] were the first ones to construct a Weyl invariant spinning membrane action. However, such action was highly *non – polynomial* complicating the quantization process . The suitable action to supersymmetrize is the one of Dolan and Tchrakian (DT) [2] without a cosmological constant and with quadratic and quartic-derivative terms. Such membrane action is basically a Skyrminion like action.

We shall write down the supersymmetrization of the polynomial DT action that is devoid of R and kinetic gravitino terms; where supersymmetry is linearly realized and the gauge algebra closes [3]. The crux of the work [3] reviewed here lies in the necessity to Weyl-covariantize the Dolan–Tchrakian action through the introduction of extra fields. These are the gauge field of dilations , b_μ , and the scalar coupling , A_0 of dimension $(length)^3$, that must appear in front of the quartic derivative terms of the DT action.

Our coventions are: Greek indices stand for three-dimensional ones; Latin indices for spacetime ones : $i, j = 0, 1, 2, \dots, D$. The signature of the $3D$ volume is $(-, +, +)$. The Dolan–Tchrakian Action for the bosonic p -brane (extendon) with vanishing cosmological constant in the case that $p = odd; p + 1 = 2n$ is :

$$L_{2n} = \sqrt{-g} g^{\mu_1 \nu_1} \dots g^{\mu_n \nu_n} \partial_{[\mu_1} X^{i_1} \dots \partial_{\mu_n]} X^{i_n} \partial_{[\nu_1} X^{j_1} \dots \partial_{\nu_n]} X^{j_n} \eta_{i_1 j_1} \dots \eta_{i_n j_n} \dots \quad (1)$$

η_{ij} is the spacetime metric and $g^{\mu\nu}$ is the world volume metric of the $2n$ hypersurface spanned by the motion of the $p - brane$. Antisymmetrization of indices is also required. Upon the algebraic elimination of the world volume

metric $g_{\mu\nu}$ from the action and after pulgging its value back into the action one recovers the Dirac–Nambu–Goto action :

$$L_{2n} = \sqrt{-\det(\partial_\mu X^i(\sigma)\partial_\nu X^j(\sigma)\eta_{ij})}. \quad (2)$$

When $p = \text{even}$, $p+1$ odd, a Lagrangian with zero cosmological constant can also be constructed, however, conformal invariance is *lost*. In the membrane's case one has : $L = L_4 + aL_2$; where a is a constant and L_2 is the standard Howe-Polyakov-Tucker quadratic actions $(\partial_\mu X^i(\sigma))^2$ and the quartic derivative terms are the Skyrmin-like terms :

$$L_4 = \sqrt{-g}g^{\mu\nu}g^{\rho\tau}\partial_{[\mu}X^i(\sigma)\partial_{\rho]}X^k(\sigma)\partial_{[\nu}X^j(\sigma)\partial_{\tau]}X^l(\sigma)\eta_{ij}\eta_{kl}. \quad (3)$$

The Supersymmetrization of the Kinetic Terms is similar to the construction of actions for the 3D Kinetic matter superconformal multiplet where supersymmetry is linearly realized and without R terms. In particular, we will show why the action is invariant under a *modified* Q -transformation , $\tilde{\delta}_Q = \delta_Q + \delta'$, which includes a compensating S and K -transformation which cancel the anomalous/spurious contributions to the ordinary Q transformations , due to the explicit breakdown of S and K -invariance of the action, which in turn, induce a breakdown of the ordinary Q supersymmetry as well.

The final action is invariant under P, D, M^{ab} transformations : translations, dilations, Lorentz. But is *not* invariant under conformal boost K and S -supersymmetry. The gauge algebra closes and the action of the modified supercovariant derivative operator $\mathcal{D}_\mu^c$ can be read from the commutator of two modified supersymmetry transformations :

$$[\tilde{\delta}_Q, \tilde{\delta}_Q](.) = \bar{\epsilon}^2 \gamma^\mu \epsilon^1 \mathcal{D}_\mu^c(.). \quad (4)$$

the modified $\mathcal{D}_\mu^c$ supercovariant derivative operator is obtained after a compensating transformation δ' is added to the standard Q -transformation to cancel the anomalous S and K -transformations of the supermultiplets used to construct the action.

The scalar and kinetic multiplet of simple conformal SG in 3D are respectively [4] :

$$\Sigma_c = (A, \chi, F). \quad T_c(\Sigma_c) = (F, \gamma^\mu D_\mu^c \chi, \triangle A). \quad (5)$$

For a supermultiplet of Weyl weight $\omega(A)$, the supercovariant derivatives and the D'Alambertian are respectively :

$$D_\mu^c A = \partial_\mu A - \frac{1}{2}\bar{\psi}_\mu \chi - \omega(A)b_\mu A. \quad (6)$$

$$D_\mu^c \chi = (D_\mu - (\omega(A) + \frac{1}{2})b_\mu)\chi - \frac{1}{2}\gamma^\mu D_\mu^c A \psi_\mu - \frac{1}{2}F\psi_\mu - \omega(A)A\phi_\mu. \quad (7)$$

$$\begin{aligned}\triangle A &= D_a^c D^{ca} A = e^{-1} \partial_\nu (e g^{\mu\nu} D_\mu^c A) + \frac{1}{2} \bar{\phi}_\mu \gamma^\mu \chi - [\omega(A) - 1] b^\mu D_\mu^c A + \\ &2\omega(A) A f_\mu^a e_a^\mu - \frac{1}{2} \bar{\psi}^\mu D_\mu^c \chi - \frac{1}{2} \bar{\psi}^\mu \gamma^\nu \psi_\nu D_\mu^c A.\end{aligned}\quad (8)$$

The generalized spin connection $\omega_\mu^{mn}(e_\mu^m, b_\mu, \psi_\mu)$ is :

$$\omega_\mu^{mn} = -\omega_\mu^{mn}(e) - \kappa_\mu^{mn}(\psi) + e_\mu^n b^m - e_\mu^m b^n. \quad (9)$$

$$\kappa_\mu^{mn} = \frac{1}{4} (\bar{\psi}_\mu \gamma^m \psi^n - \bar{\psi}_\mu \gamma^n \psi^m + \bar{\psi}^m \gamma_\mu \psi^n). \quad (10)$$

The gravitino field strength is :

$$\phi_\mu = \frac{1}{4} \sigma^{\lambda\sigma} \gamma_\mu S_{\sigma\lambda}. \quad (11)$$

$$S_{\mu\nu} = (D_\nu + \frac{1}{2} b_\nu) \psi_\mu - \mu \leftrightarrow \nu. \quad (12)$$

$$e^{a\mu} f_{a\mu} = -\frac{1}{8} R(e, \omega) - \frac{1}{4} \bar{\psi}_\mu \sigma^{\mu\nu} \phi_\nu. \quad (13)$$

the variations of the gauge and matter fields under D (dilatations) , and Q, K, S , transformations is :

$$\delta_D e_\mu^m = -\lambda e_\mu^m. \quad \delta_D A = \frac{1}{2} \lambda A. \quad \delta_D F = \frac{3}{2} \lambda F. \quad (14)$$

In general, a Weyl supermultiplet of weight ω transforms under Weyl scalings as :

$$\delta_D A = \omega \lambda A. \quad \delta_D \chi = (\omega + \frac{1}{2}) \lambda \chi. \quad \delta_D F = (\omega + 1) \lambda F. \quad (15)$$

where λ is the Weyl scaling's infinitesimal parameter. Under Q -supersymmetry :

$$\delta_Q A = \bar{\epsilon} \chi. \quad \delta_Q \chi = F \epsilon + \gamma^\mu D_\mu^c A \epsilon. \quad \delta_Q F = \bar{\epsilon} \gamma^\mu D_\mu^c \chi. \quad (16)$$

$$\delta_Q e_\mu^m = \bar{\epsilon} \gamma^m \psi_\mu. \quad \delta_Q \psi_\mu = 2(D_\mu + \frac{1}{2} b_\mu) \epsilon. \quad \delta_Q b_\mu = \phi_\mu. \quad (17)$$

Under S -supersymmetry :

$$\delta_S e_\mu^m = 0. \quad \delta_S \psi_\mu = -\gamma_\mu \epsilon_s. \quad \delta_S b_\mu = -\frac{1}{2} \psi_\mu \epsilon_s. \quad (18)$$

$$\delta_S \omega_\mu^{mn} = -\bar{\epsilon}_s \psi_\mu. \quad \delta_S A = 0. \quad \delta_S \chi = \omega(A) A \epsilon_s. \quad \delta_S F = (1 - \frac{1}{2} \omega(A)) \bar{\chi} \epsilon_s. \quad (19)$$

The canonical scalar supermultiplet is *inert* under conformal boosts K and so are the e_μ^m, ψ_μ . The b_μ is *not* inert :

$$\delta_K b_\mu = -2\xi_K^m e_{m\mu}. \quad \delta_K \omega_\mu^{mn} = 2(\xi_K^m e_\mu^n - \xi_K^n e_\mu^m). \quad (20)$$

A superconformally invariant action for the kinetic terms in $D = 3$ requires to include *only* the following kinetic multiplet that is fully superconformally invariant : $\Sigma_C \otimes T_C(\Sigma_C)$. The superconformal invariant action was given in [4].

However, in this work we shall use the different combination of supermultiplets

$$\Sigma_C^i \otimes T_C(\Sigma_C^j) + T_C(\Sigma_C^i) \otimes \Sigma_C^j - T_C(\Sigma_C^i \otimes \Sigma_C^j). \quad (21)$$

combination which happens to be the correct one to dispense of the R and kinetic gravitino terms. However this combination *breaks* explicitly the S and conformal boosts. In doing so, it also will break the Q -invariance of any action constructed based on such “anomalous” kinetic multiplet [3].

The explicit components of the suitable combination of multiplets which removes the R and gravitino kinetic terms and breaks K -symmetry and S -supersymmetry ,

$$\Sigma_C^i \otimes T_C(\Sigma_C^j) + T_C(\Sigma_C^i) \otimes \Sigma_C^j - T_C(\Sigma_C^i \otimes \Sigma_C^j). \quad (22)$$

are :

$$A_{ij} = \bar{\chi}_i \chi_j. \quad \chi_{ij} = F_i \chi_j + F_j \chi_i + A_i \gamma^\mu D_\mu^c(\omega = \frac{1}{2}) \chi_j + A_j \gamma^\mu D_\mu^c(\omega = \frac{1}{2}) \chi_i -$$

$$\gamma^\mu D_\mu^c(\omega = 1) [A_i \chi_j + A_j \chi_i].$$

$$F_{ij} = A_i \Delta(\omega = \frac{1}{2}) A_j + A_j \Delta(\omega = \frac{1}{2}) A_i + 2F_i F_j - \bar{\chi}_i \gamma^\mu D_\mu^c(\omega = \frac{1}{2}) \chi_j -$$

$$\bar{\chi}_j \gamma^\mu D_\mu^c(\omega = \frac{1}{2}) \chi_i - \Delta(\omega = 1) [A_i A_j]. \quad (23)$$

The explicit expression for the multiplet components was given in [3]. We can explicitly verify [3] that there is no gravitino kinetic term and no curvature scalar terms, as expected. There is an explicit presence of the b_μ terms, which signals an explicit breaking of conformal boosts. Therefore, eliminating the R and gravitino kinetic terms is *not* compatible with S -supersymmetry nor K -symmetry.

Because the component $T_C(\Sigma_C \otimes \Sigma_C)$ does not have the correct $S; K$ transformations laws, the components of the latter supermultiplet ($A_{ij}; \chi_{ij}; F_{ij}$) *do not* longer transform properly under (ordinary) Q transformations because the supercovariant derivatives acting on the A_{ij}, χ_{ij} acquire anomalous/spurious

terms due to the explicit S and K -breaking pieces [3]. To correct this problem one needs to modify the Q supersymmetry by adding compensating S, K transformations as shown below.

The supersymmetric quadratic derivative terms (kinetic action) of the action are obtained by plugging in directly the components $A_{ij}, \chi_{ij}, F_{ij}$ into the following expression while contracting the spacetime indices with η_{ij} :

$$\mathcal{L}_2 = e\eta^{ij}[F_{ij} + \frac{1}{2}\bar{\psi}_\mu\gamma^\mu\chi_{ij} + \frac{1}{2}A_{ij}\bar{\psi}_\mu\sigma^{\mu\nu}\psi_\nu]. \quad (24)$$

After using a modified Q -variation of the action the spurious variations of the action, resulting from the S and K -symmetry breaking pieces in the definitions of supercovariant derivatives, will *cancel* and the action is invariant , up to total derivatives. [3]. Using the modified derivatives, resulting from the cancellation of the anomalous S, K variations of the the supermultiplet, the action obtained from eq-(24) will be invariant under the following *modified* Q -transformations :

$$\tilde{\delta}_Q(A_{ij}) = \delta_Q^c(A_{ij}) = \bar{\epsilon}\chi_{ij}, \quad \tilde{\delta}_Q(\chi_{ij}) = F_{ij}\epsilon + \gamma^\mu\mathcal{D}_\mu^c(A_{ij})\epsilon. \quad (25)$$

$$\tilde{\delta}_Q F_{ij} = \bar{\epsilon}\gamma^\mu\mathcal{D}_\mu^c(\chi_{ij}). \quad (26)$$

where now is essential to use the modified supercovariant derivatives $\mathcal{D}_\mu^c$ associated with the modified Q -transformations laws in order to cancel the anomalous contributions stemming from the S and K -breaking pieces, since the latter anomalous contributions to the Q -variations are contained in the *ordinary* derivatives $D_\mu^c(A_{ij}); D_\mu^c(\chi_{ij})$.

Under ordinary Q -variations the Lagrangian density will acquire spurious pieces [3] : $(\Delta L)|_{spurious} = \delta_Q[eL]|_{spurious}$. What the modified $\tilde{\delta}_Q\mathcal{S} = [\delta_Q + \delta']\mathcal{S}$ achieves is to have $\delta'\mathcal{S} = -\Delta|_{spurious}$ so that the net *sum* of the anomalous Q, K, S contributions cancels out giving zero at the end ; i.e one has the correct $Q + K + S$ sum rule.

The modified covariant derivatives, appearing in the modified Q transformation laws of the supermultiplet of Weyl weight $\omega = 2$, used to construct the action , have now the appropriate form to ensure invariance of the actions under the modified $\tilde{Q}$ transformations :

$$\mathcal{D}_\mu^c(A_{ij}) = \partial_\mu A_{ij} - \frac{1}{2}\bar{\psi}_\mu\chi_{ij} - 2b_\mu A_{ij}. \quad (27)$$

$$\mathcal{D}_\mu^c\chi_{ij} = (D_\mu - (2 + \frac{1}{2})b_\mu)\chi_{ij} - \frac{1}{2}\gamma^\nu\mathcal{D}_\nu^c(A_{ij})\psi_\mu - \frac{1}{2}F_{ij}\psi_\mu - 2A_{ij}\phi_\mu. \quad (28)$$

Since $\tilde{\delta}_Q\mathcal{S} = 0$, one has found the correct $Q + K + S$ sum rule for the modified Q transformation. Finally, we don't have R terms, nor the kinetic terms for the gravitino ; i.e there is no f_μ^m term. Q -supersymmetry is linearly

realized after the elimination of F^i . No constraints arise after eliminating the auxiliary fields F^i [3].

The Supersymmetrization of the Quartic Derivative Terms proceeds in analogous fashion. Let us introduce the coupling-function supermultiplet, $\Sigma_0 = (A_0; \chi_0; F_0)$ whose Weyl weight is equal to -3 so that the tensor product of Σ_0 with the following multiplet, to be defined below, has a conformal weight, $\omega = 2$ as it is required in order to have Weyl invariant actions. The following multiplet is the adequate one to build the required supersymmetrization of the quartic derivative terms :

$$K_{\mu\nu\rho\tau}^{ijkl} = K(\Sigma_\mu^i; \Sigma_\nu^j) \otimes T[K(\Sigma_\rho^k; \Sigma_\tau^l)] + (ij \leftrightarrow kl) \text{ and } (\mu\nu \leftrightarrow \rho\tau). \quad (29)$$

$$T[K(\Sigma_\mu^i; \Sigma_\nu^j) \otimes K(\Sigma_\rho^k; \Sigma_\tau^l)]. \quad (30)$$

$$K(\Sigma, \Sigma) = \Sigma_C^i \otimes T_C(\Sigma_C^j) + T_C(\Sigma_C^i) \otimes \Sigma_C^j \equiv (A_{ij}^K; \chi_{ij}^K; F_{ij}^K). \quad (31)$$

This combination of these supermultiplets is the adequate one to retrieve the DT action at the bosonic level and also ensures that the R terms do cancel from the final answer. This is a similar case like the construction of the quadratic terms which was devoid of R and kinetic gravitino terms. A similar calculation yields the components of the supersymmetric-quartic-derivative terms:

$$\begin{aligned} A_{ijkl} &= \bar{\chi}_{ij}^K \chi_{kl}^K, \quad \chi_{ijkl} = F_{ij}^K \chi_{kl}^K + F_{kl}^K \chi_{ij}^K + \\ &A_{ij}^K \gamma^\mu D_\mu^c(\omega = 2) \chi_{kl}^K + A_{kl}^K \gamma^\mu D_\mu^c(\omega = 2) \chi_{ij}^K - \\ &\gamma^\mu D_\mu^c(\omega = 4) [A_{ij}^K \chi_{kl}^K + A_{kl}^K \chi_{ij}^K]. \end{aligned} \quad (32)$$

$$F_{ijkl} = A_{ij}^K \Delta(\omega = 2) A_{kl}^K + A_{kl}^K \Delta(\omega = 2) A_{ij}^K + 2F_{ij}^K F_{kl}^K -$$

$$\bar{\chi}_{ij}^K \gamma^\mu D_\mu^c(\omega = 2) \chi_{kl}^K - \bar{\chi}_{kl}^K \gamma^\mu D_\mu^c(\omega = 2) \chi_{ij}^K - \Delta(\omega = 4) [A_{ij}^K A_{kl}^K]. \quad (33)$$

where we have used the abbreviations A_{ij}^K, χ_{ij}^K and F_{ij}^K given by the well behaved kinetic multiplet.

Despite the fact that the defining quartic multiplet is devoid of curvature terms we are once again faced with the anomalous Q transformations of the action due to the spurious S, K -variations. To cure these anomalous variations of the action we proceed exactly as before by using a modified Q transformations that will cancel the spurious variations.

The complete Q supersymmetric extension of L_4 requires adding terms which result as permutations of $ijkl \leftrightarrow ilkj \leftrightarrow kjil \leftrightarrow klij$ keeping $\eta_{ij}\eta_{kl}$ fixed. The action corresponding to the quartic derivative terms is :

$$\begin{aligned} \mathcal{L}_4 = e\eta_{ij}\eta_{kl} [A_0 F^{ijkl} + F_0 A^{ijkl} - \bar{\chi}_0 \chi^{ijkl} + \frac{1}{2} \bar{\psi}_\mu \gamma^\mu (A_0 \chi^{ijkl} + \chi_0 A^{ijkl}) + \\ \frac{1}{2} A_0 A^{ijkl} \bar{\psi}_\mu \sigma^{\mu\nu} \psi_\nu]. \end{aligned} \quad (34)$$

The composite fields A_{ijkl} , χ_{ijkl} and F_{ijkl} are given in terms of the $A_{ij}^K, \chi_{ij}^K, F_{ij}$. The action will be invariant under the *modified* $\mathcal{Q}$ -supersymmetry transformations laws, which have the same structure as the variations of the quadratic derivatives multiplet :

$$\tilde{\delta}_Q \mathcal{A}_{ijkl} = \bar{\epsilon} \chi_{ijkl}. \quad \tilde{\delta}_Q \chi_{ijkl} = F_{ijkl} \epsilon + \gamma^\mu \mathcal{D}_\mu^c \mathcal{A}_{ijkl} \epsilon. \quad \tilde{\delta}_Q F_{ijkl} = \bar{\epsilon} \gamma^\mu \mathcal{D}_\mu^c \chi_{ijkl}. \quad (35)$$

where

$$\mathcal{D}_\mu^c (A_{ijkl}) = \partial_\mu A_{ijkl} - \frac{1}{2} \bar{\psi}_\mu \chi_{ijkl} - 2b_\mu A_{ijkl}. \quad (36)$$

and similarly for :

$$\mathcal{D}_\mu^c \chi_{ijkl} = (D_\mu - (2 + \frac{1}{2})b_\mu) \chi_{ijkl} - \frac{1}{2} \gamma^\nu D_\nu^c (A_{ijkl}) \psi_\mu - \frac{1}{2} F_{ijkl} \psi_\mu - 2A_{ijkl} \phi_\mu. \quad (36)$$

Concluding, we have presented all the steps necessary to construct a satisfactory Weyl invariant spinning membrane action that is : (i) Polynomial in the fields (ii) devoid of curvature terms (iii) where supersymmetry is linearly realized . (iv) Upon setting the fermions to zero, and eliminating the auxiliary fields, it yields the Weyl-covariant extension of the Dolan-Tchrakian action for the membrane.

Acknowledgements

We thank M. Bowers and J. Mahecha for their assistance.

References

- [1] U.Lindstrom, M. Rocek, Phys. Letters **B 218** (1988) 207;
- [2] B.P.Dolan, D.H.Tchrakian, Physics Letters **B 198** (1987) 447;
- [3] C. Castro, “ Remarks on the existence of Spinning Membrane Actions “ hep-th/0007031;
- [4] T.Uematsu, Z.Physics **C 29** (1985) 143-146 and **C32** (1986) 33-42;
- [5]. M. Duff, S. Deser, C. Isham , Nuc. Phys. **B 114** (1976) 29;
- [6] . E. Guendelman : “ Superextendons with a modified Measure “ hep-th/0006079.